

Lecture 4

Shape of distribution and Measures of Central Tendency

Review From Friday 1/19

3 features of a distribution that we are interested in:

- Shape
- Center
- Spread or variability

Graphs of data are a good way summarize patterns in data

Graphs for qualitative data are

- Bar graphs, pie charts

Graphs for quantitative data are:


- Stem plot, dot plot, histogram

Practice: Histogram

- $X = \{-1.49, -0.65, -0.6, -0.54, -0.45, 0.01, 0.17, 0.27, 0.51, 1.34\}$


Construct a histogram using $K = 4$ bins/intervals:

Shape of a distribution




- Bimodal distributions can arise when
 - A population is polarized on a controversial issue
 - When observations come from two different sub-populations

Histogram of Eruption Waiting Times




Democrats and Republicans More Ideologically Divided than in the Past

Distribution of Democrats and Republicans on a 10-item scale of political values




Source: 2014 Political Polarization in the American Public
 Notes: Ideological consistency based on a scale of 10 political values questions (see Appendix A). The blue area in this chart represents the ideological distribution of Democrats; the red area of Republicans. The overlap of these two distributions is shaded purple. Republicans include Republican-leaning independents; Democrats include Democratic-leaning independents (see Appendix B).

PEW RESEARCH CENTER




- Skewed distributions occur when there is a strict boundary on the possible values of a variable

- Consider the following histogram of median housing prices in California from the 1990 national census



- **Outliers** are extreme values that fall far away from the midpoint of the data
- Consider the following histogram of the fuel efficiency of cars from 1990 - 2008



Measures of Central Tendency

- The (arithmetic) **mean** is the average value of a set of observations
it measures the center of mass of a distribution (the balancing point)

$$\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \sum_{i=1}^n \frac{x_i}{n}$$

- the mean is usually not equal to any of the values observed in the sample
- The mean is highly influenced by **outliers** - observations that take on extreme values relative to the distribution

Practice: Calculate The Mean

- $X = \{1, 3, 5, 5, 6, 7, 7, 8\}$

Measures of Central Tendency

- The **median** is the middle value of a set of observations


How to compute the median:

1. Compute the median by first ordering the observations from smallest value to largest value and choose the number in the middle
2. If the n is odd the median is the middle number
 - If n is even the median is the sum of the two middle values divided by 2

Practice: Calculate the Median


- $X = \{1, 3, 5, 5, 6, 7, 7, 8\}$

Data: 3 3 3 3 3 3 3 4 4 4 4 4 4 4 4 4 5 5 5 5 5 5 5 5 5 5
 5 5 5 5 5 5 5 5 6 6 6 6 6 6 6 6 6 6 7 7 7 7 7 7 7
 7 8 8 8 8 8 9 9 9 9 9 9 10 10 10 10 10 10 10 11 11 11 11 11 12
 12 12 12 12 12 12 12 13 13 13 14 14 14 14 14 14 15 15 15 16 16 16 16 17 17 17 18
 18 20 20 20 20 20



$$\bar{x} = 9.2$$

The mean is the center of gravity







$$median = \frac{8 + 8}{2} = 8$$

The median is the middle value

Mean and median treat outliers differently

- $X = \{1, 3, 5, 5, 6, 7, 7, 8, 32\}$



Alternative formulas for the mean

- We can also express the mean in terms of the frequency F or the relative frequency RF

$$\bar{x} = \frac{1}{n} \sum_x x F(x) \quad \text{or} \quad \bar{x} = \sum_x x RF(x)$$

Where the sum is over all distinct values of the variable x

Example: Computing the mean from a frequency table

$X = \{1, 3, 5, 5, 6, 7, 7, 8\}$

| X | Freq. | Rel. Freq |
|---|-------|-----------|
| 1 | 1 | 0.125 |
| 3 | 1 | 0.125 |
| 5 | 2 | 0.250 |
| 6 | 1 | 0.125 |
| 7 | 2 | 0.250 |
| 8 | 1 | 0.125 |

The mode

- The **mode** is the value with the largest relative frequency (i.e the value that occurs most often)
 - Can be used with categorical data (mean and median cannot)
 - e.g the most frequent category
 - It may not be unique if two or more values have the same frequency
 - **Caution** for quantitative data, the mode may not anywhere near the center of the distribution.

Ex.)

Data = 1,1,4,5,6

Mode = 1

Data = 1,1,4,5,6,6

Mode 1, 6

Practice:

- Roll a six-sided die $n = 10$ times and record the number rolled each time
- Data = 1,2,3,3,4,4,4,5,6,6

| x | $f(x)$ | $rf(x)$ |
|-----|--------|---------|
| 1 | 1 | 0.1 |
| 2 | 1 | 0.1 |
| 3 | 2 | 0.2 |
| 4 | 3 | 0.3 |
| 5 | 1 | 0.1 |
| 6 | 2 | 0.2 |

Compute the **mean** using all 3 equations:

Compute the **median**

Compute the **mode**